

PATENT APPLICATION SERIAL NO.

08/238080

U.S. DEPARTMENT OF COMMERCE
PATENT AND TRADEMARK OFFICE
FEE RECORD SHEET

08/238080

PTO-1556
(5/87)

MM20121 05/18/94 08238080

01-0528 020 101

872.00CH 2583511



08/238,080

ABSTRACT

A method of assay for target polynucleotides includes steps of isolating target polynucleotides from extraneous non-target polynucleotides, debris, and impurities and amplifying the target polynucleotide.

5

10

15

20

25

30

35

08238080-050394



TARGET AND BACKGROUND CAPTURE METHODS WITH AMPLIFICATION FOR AFFINITY ASSAYS

08/238080

malt
08/05/95

BACKGROUND OF THE INVENTION

This application is a continuation-in-part of U.S.S.N. 922,155, entitled TARGET AND BACKGROUND CAPTURE METHODS FOR AFFINITY ASSAYS, filed October 1986, and hereby incorporated by reference.

The present invention pertains to methods, reagents, compositions, kits, and instruments for use in capturing target molecules. In particular, the present invention relates to methods, reagents, compositions, and kits for capturing deoxyribonucleic acid (DNA) or ribonucleic acid (RNA) from clinical samples. Embodiments of the present invention provide methods for rapid, sensitive detection of nucleic acid targets in clinical samples adaptable to non-radioactive labeling techniques and automation.

The following definitions are provided to facilitate an understanding of the present invention. The term "biological binding pair" as used in the present application refers to any pair of molecules which exhibit natural affinity or binding capacity. For the purposes of the present application, the term "ligand" will refer to one molecule of the biological binding pair and the term "antiligand" or "receptor" will refer to the opposite molecule of the biological binding pair. For example, without limitation, embodiments of the present invention have applications in nucleic acid hybridization assays where the biological binding pair includes two complementary strands of polynucleic acid. One of the strands is designated the ligand and the other strand is designated the antiligand. However, the biological binding pair may include antigens and antibodies, drugs, and drug receptor sites and enzymes and enzyme substrates.

The term "probe" refers to a ligand of known qualities capable of selectively binding to a target antili-

2

1

10

15

Figure 1 consists of 12 micrographs arranged vertically, illustrating the stages of chick embryo development. The stages are labeled as follows from top to bottom:

- 1. Fertilized egg (single cell)
- 2. 2-cell stage
- 3. 4-cell stage
- 4. 8-cell stage
- 5. Morula stage
- 6. Gastrula stage
- 7. Head stage
- 8. Heart stage
- 9. Somite stage
- 10. Limb stage
- 11. Hatching stage
- 12. Fully hatched chick

25

35

can be made enzymatically with DNA or RNA polymerases or transcriptases.

Genetic information is stored in living cells in threadlike molecules of DNA. In vivo, the DNA molecule is a double helix, each strand of which is a chain of nucleotides. Each nucleotide is characterized by one of four bases: adenine (A), guanine (G), thymine (T), and cytosine (C). The bases are complementary in the sense that, due to the orientation of functional groups, certain base pairs attract and bond to each other through hydrogen bonding. Adenine in one strand of DNA pairs with thymine in an opposing complementary strand. Guanine in one strand of DNA pairs with cytosine in an opposing complementary strand. In RNA, the thymine base is replaced by uracil (U) which pairs with adenine in an opposing complementary strand.

DNA consists of covalently linked chains of deoxyribonucleotides and RNA consists of covalently linked chains of ribonucleotides. The genetic code of a living organism is carried upon the DNA strand in the sequence of the base pairs.

Each nucleic acid is linked by a phosphodiester bridge between the five prime hydroxyl group of the sugar of one nucleotide and the three prime hydroxyl group of the sugar of an adjacent nucleotide. Each linear strand of naturally occurring DNA or RNA has one terminal end having a free five prime hydroxyl group and another terminal end having a three prime hydroxyl group. The terminal ends of polynucleotides are often referred to as being five prime termini or three prime termini in reference to the respective free hydroxyl group. Complementary strands of DNA and RNA form antiparallel complexes in which the three prime terminal end of one strand is oriented to the five prime terminal end of the opposing strand.

Nucleic acid hybridization assays are based on the tendency of two nucleic acid strands to pair at comple-

08238080-1510394
156051-08082380

5

10

25

30

35

5

[illegible]

5

10

20

30

35

external energy source and may be grouped into categories dependent upon the source of the exciting energy, including: radioluminescent labels deriving energy from high energy particles; chemiluminescent labels which obtain energy from chemical reactions; bioluminescent labels wherein the exciting energy is applied in a biological system; and photoluminescent or fluorescent labels which are excitable by units of electromagnetic radiation (photons) of infrared, visual or ultraviolet light. See, generally, Smith et al., Ann. Clin. Biochem., 18: 253, 274 (1981).

Nonisotopic assay techniques employing labels excitable by nonradioactive energy sources avoid the health hazards and licensing problems encountered with radioisotopic label assay techniques. Moreover, nonisotopic assay techniques hold promise for rapid detection avoiding the long exposure time associated with the use of X-ray film.

However, nonisotopic assays have not conveyed the sensitivity or specificity to assay procedures necessary to be considered reliable. In luminescent assays, the presence of proteins and other molecules carried in biological samples may cause scattering of the exciting light or may absorb light in the spectrum of emission of the luminescent label, resulting in a quenching of the luminescent probe.

In enzymatic assays, the presence of proteins and other molecules carried in biological samples may interfere with the activity of the enzyme.

Similarly, in colorimetric assays, the change in color may not be detectable over proteins and other materials carried in biological samples.

Embodiments of the present invention are concerned with target and background capture on supports and on retrievable supports including magnetic particles. Magnetic particles have been suggested as supports for the synthesis of organic compounds, including oligomers such

as DNA, RNA, polypeptides, and other multiunit molecules that have a defined sequences. See, for example, European Patent Application No. 83112493.8 to Steven A. Benner and Genetics Institute. However, magnetic particles have not been suggested as retrievable supports for target capture and background removal.

Other utilization of magnetic particles has included magnetic fluids in the blood, R. Neubauer, IEEE transactions on magnetics MAG-9, 445 (1973); attachment of functional group for separation of biomolecules, U.S. Patent No. 3,970,518 to I. Giaver; labelling of cell-surface receptors, S. Margel et al., Jour. Imm. Meth. 28:341-53 (1979); attachment to drugs for magnetic targeting during therapeutic, A. Senyei et al., J., App. Phys., 49 (6): 3578 (1978), K. Wieder et al., Pro. Soc. of Exp. Bio. Med., 58:141 (1978), K. Mosbach and U. Shroeder, FEBS letters 102:112 (1979); selective separation of viruses, bacteria, and other cells, R. Molday et al., Nature 268:438 (1977); and incorporation of magnetic particles as support in gel affinity chromatography for biological polymers, K. Mosbach and L. Anderson, Nature 270:359 (1977), which are incorporated herein by reference.

The use of a two probe system to effect target capture on conventional non-retrievable supports has been suggested in an article authored by Ann-Christine Syuänen, Matti Laaksonen and Hans Söderlund entitled "Fast Quantification of Nucleic Acid Hybrids by Affinity-Based Hybrid Collection;" Nucleic Acids Research, 14(12): 5037 (1986).

SUMMARY OF THE INVENTION

It is an object of the present invention to provide methods, reagents, compositions, kits, and instrumentation for performing assays for target molecules of interest. Other objects will be presented hereinafter. For convenience, without limitation embodiments of the present invention can be grouped into areas of target

465050-0308E280

8

5

15

25

30

35

T

The retrievable support is capable of substantially homogenous dispersion within a sample medium. The probe moiety may be associated to the retrievable support, by way of example, by covalent binding of the probe moiety to the retrievable support, by affinity association, hydrogen bonding, or nonspecific association.

The support may take many forms including, by way of example, nitrocellulose reduced to particulate form and retrievable upon passing the sample medium containing the support through a sieve; nitrocellulose or the materials impregnated with magnetic particles or the like, allowing the nitrocellulose to migrate within the sample medium upon the application of a magnetic field; beads or particles which may be filtered or exhibit electromagnetic properties; and polystyrene beads which partition to the surface of an aqueous medium.

A preferred embodiment of the present invention includes a retrievable support comprising magnetic beads characterized in their ability to be substantially homogeneously dispersed in a sample medium. Preferably, the magnetic beads carry primary amine or carboxyl functional groups which facilitate covalent binding or association of a probe entity to the magnetic support particles. Preferably, the magnetic support beads are single domain magnets and are super paramagnetic exhibiting no residual magnetism. The first probe includes a probe ligand moiety capable of specifically binding to antiligand under binding conditions. The retrievable support is capable of substantially homogeneous dispersion within the sample media and includes at least one antiligand moiety capable of binding to ligand under binding conditions to form a target-probe support complex. Next, the retrievable support and sample medium are separated to allow the sample medium to be processed further.

Embodiments of the invention are suitable for capturing target molecules from a clinical sample medium containing extraneous material. The order of contacting

44-0000-00000000

the sample medium with probe or retrievable support is a matter of choice. However, the choice may be influenced by the kinetics of binding between the probe and target on one hand, and between the probe ligand and support antiligand on the other.

As applied to polynucleotide target molecules and homopolymer ligands and antiligands, the homopolymer ligand and antiligand binding is generally faster than probe binding to target. Probe binding to the target is sterically impaired after the probe ligand is bound to the support antiligand. A preferred embodiment includes contacting the sample medium with the reagent and bringing the mixture to hybridization conditions. Next, the retrievable support is dispersed in the reagent and sample medium allowing the formation of a target-probe complex in advance of the formation of probe support complexes.

A further embodiment of the present invention features a multiple probe system.

Preferably the method includes a reagent including a first probe as previously described and at least one second probe capable of binding to the target molecule and having label moieties capable of detection. The second probe is capable of forming a target (first and second) probe-support complex. The step of separating the retrievable support from the sample medium not only removes extraneous material from the target-(first and second) probe-support complex, but also separates any second probe which is not bound to the target. Second probe unbound to target contributes to background noises, false signals indicating the presence of target.

Further processing may include release of the target (first and second) probe complex from the retrievable support into a second medium and rebinding of the target (first and second) probe complex to new support. The first retrievable support may carry nonspecifically bound materials which can interfere with assay procedures.

Thus, after the release of the target-probe complex from the retrievable support and the retrievable supports removal, a second support having an antiligand moiety capable of binding to the probe ligand can be brought into contact with the target-probe complex under binding conditions to effect a further cycle of target-probe binding or capture for further purification and concentration of target-probe complex.

Further processing may include background capture. A further embodiment of the present invention includes a method wherein the second probe has a second ligand moiety. The method further includes a background support having a second antiligand moiety. The second ligand moiety and second antiligand moiety are capable of stably binding under binding conditions only when the second probe is unbound to the target molecule. The method further includes the step of contacting a medium potentially containing second probe unbound to target with a background support under binding conditions. Next, the background support is separated from the medium to remove unbound second probe reducing background noise.

The term "background support" is used in the conventional sense to include filters and membranes as well as retrievable supports. Binding to the background support does not need to be releasible.

A preferred retrievable support includes, by way of example without limitation, particles, grains, beads, or filaments capable of dispersion within and separation from a medium. Methods of separation include by way of example, without limitation, of filtration, centrifugation, precipitation, surface floatation, settling, or the introduction of electromagnetic fields.

The present method can be applied to polynucleotide target molecules. Preferably, the first and second probes bind quickly to a polynucleotide target "in solution" as opposed to the situation where either the target or probe is immobilized.

452951-0808E230

The retrievable support, capable of substantial dispersion within a solution, permits interactions between the retrievable support and probes which mimic "in solution" hybridization. In solution, hybridization can be completed in approximately 3-15 minutes. The rapid hybridizations and simplicity of the present methods permit automation. The present method allows nucleic acid sequences contained in clinical samples to be separated from extraneous material allowing the methods to be applied to nonisotopic labeling techniques.

An embodiment of the present method where the target molecule is a polynucleotide, includes contacting a sample medium with reagent under binding conditions. The reagent includes at least one first polynucleotide probe and at least one second polynucleotide probe. The first probe is capable of forming a complex with the target molecule and has a first homopolymer ligand moiety. The second probe is capable of forming a complex with the target molecule in addition to the first probe. The second probe includes a label moiety which has a second homopolymer ligand moiety which is different than the first homopolymer ligand of the first probe. Next, the reagent and sample medium are contacted with a background support and a target capture support. The background support includes at least one second homopolymer antiligand moiety capable of binding to the second homopolymer ligand moiety of the second probe when said second probe is unbound to target. The target capture support includes at least one first homopolymer moiety capable of binding to the first homopolymer ligand moiety of the first probe. The background support and the target capture support remove background noise and the target capture support further concentrates the target-(first and second) probe complex for further processing and separates the target-(first and second) probe complex from cellular debris. Further processing includes the detection of the label moiety indicative of the presence of

142-0331-0208E2B0

the target molecule.

Turning now more specifically to embodiments of the invention pertaining to background capture, one embodiment includes a method wherein probe and target are allowed to form a complex. Next, uncomplexed probe is brought into contact with a support under binding conditions. The support is capable of selectively binding unbound probe. Next, the support is separated from the probe-target complex.

A still further embodiment of the present invention includes a method of separating a plurality of target molecules for further processing.

One embodiment includes the sequential addition and removal of probes specific to target molecules on a plurality of supports. A further embodiment includes a method which includes contacting a sample with a first series probe and capturing the target and probe on a plurality of supports. The first series probe includes a ligand capable of association with the support. The first probe series includes probes for all plurality targets which are capable of binding to supports specific for each target molecule. The supports are capable of being separated from each other, the separation of which results in individual types of target molecules being isolated with the support.

A further embodiment of the present invention includes a reagent composition. The reagent composition includes a first probe and a second probe. The first probe is capable of forming a complex with a target molecule and includes a probe ligand moiety capable of specifically binding to antiligand under binding conditions. The second probe is capable of forming a complex with the target molecule and includes a label moiety capable of detection. The reagent composition can be used to capture and detect the target in a sample medium when used with a retrievable support having antiligand moieties.

5 A further embodiment of the present reagent composition includes a second probe having a second ligand moiety capable of stably binding to an antiligand only in the situation where the second probe is unbound to the target molecule. The reagent composition allows background noise to be reduced by contacting sample potentially containing an unbound second probe with a background support having a second antiligand moiety.

10 A further embodiment includes a support capable of substantially homogeneous dispersion in a sample medium having oligonucleotide antiligands adapted for binding to oligonucleotide ligands on probes.

15 A preferred embodiment of the support includes, by way of example, particles, grains, filaments, and beads capable of separation. Means of separation include, by way of example without limitation, precipitation, settling, floatation, filtration, centrifugation, and electromagnetism.

20 A preferred embodiment includes polystyrene beads, between 10-100 microns in diameter, which are capable of substantially homogeneous dispersion and separation from a medium by filtration or floatation. Another preferred embodiment includes ferromagnetic beads. A ferromagnetic bead marketed under the trademark BIO-MAG is capable of
25 substantially homogeneous dispersion in an aqueous medium and can be retrieved or immobilized by an electromagnetic field. The ferromagnetic bead includes an iron core which is coated with an amine reactive covering. The beads are generally spherical and have a diameter of one
30 micron. The polystyrene and ferromagnetic beads are treated to include antiligand moieties.

A further embodiment of the present invention includes a kit for performing assays for target molecules which are part of a biological binding pair. In the case
35 where the target is a polynucleotide having a specific base sequence, the kit includes a reagent wherein the reagent includes a first polynucleotide probe and a

16E0391:0909E280

15

[illegible]

20

25

[illegible]

5

10

15

20

25

35

amplifying the target polynucleotide, to form an amplification product, and measuring or detecting the amplified target polynucleotide. This method advantageously can be used in conjunction with the target capture and background capture steps described above.

Brief Description of the Drawings

Figs. 1-3 are flow diagrams illustrating steps, apparatus, and reagents used in methods of the invention. The term "Figure 1" refers collectively to Figure 1a and Figure 1b. Similarly, the term "Figure 2" refers collectively to Figure 2a and Figure 2b.

Figs. 4-6 are diagrammatic representations of capture amplification methods of the invention.

Fig. 7 is a diagram illustrating features of an apparatus made in accordance with one embodiment of the present invention.

Fig. 8 is a diagrammatic representation of a genetic construction used in the invention.

Detailed Description

Turning now to the drawings, which by way of illustration depict preferred embodiments of the present invention, and in particular Figure 1, a method of procedure, with necessary reagent compositions, is illustrated in schematic form for an assay for target polynucleotide strands. Conventional assay techniques include many target strands, and many probe strands would be used to perform an assay. However, for the simplicity to further an understanding of the invention, the illustration depicts only limited numbers of probes, support entities, and targets. Figure 1 features a method utilizing retrievable supports.

Step 1 of the assay illustrated in Figure 1 begins with a clinical sample which, by way of illustration, contains cells. The cells potentially carry target nucleic acid, either DNA or RNA, having a base sequence

18

20

25

20

35

binding of the nucleic acid to filters or membranes and washing cellular debris and impurities from the filter or membrane. However, in practice, some cellular debris and some impurities, as well as nontarget nucleic acid, are nonspecifically bound to the filter or membrane and are not removed by washes.

An embodiment of the present invention, as illustrated in Step 1, includes contacting the sample potentially carrying target nucleic acid with a retrievable support in association with a probe moiety. The retrievable support is capable of substantially homogenous dispersion within a sample medium. The probe moiety may be associated to the retrievable support, by way of example, by covalent binding of the probe moiety to the retrievable support, by affinity association, hydrogen binding, or nonspecific association.

The support may take many forms including, by way of example, nitrocellulose reduced to particulate form and retrievable upon passing the sample medium containing the support through a sieve; nitrocellulose or the materials impregnated with magnetic particles or the like, allowing the nitrocellulose to migrate within the sample medium upon the application of a magnetic field; beads or particles which may be filtered or exhibit electromagnetic properties; and polystyrene beads which partition to the surface of an aqueous medium.

A preferred embodiment of the present invention includes a retrievable support comprising magnetic beads characterized in their ability to be substantially homogeneously dispersed in a sample medium. Preferably, the magnetic beads contain primary amine functional groups which facilitate covalent binding or association of a probe entity to the magnetic support particles. Preferably, the magnetic support beads are single domain magnets and are super paramagnetic exhibiting no residual magnetism.

08230080 050394
+65050-08082300

5 The particles or beads may be comprised of magnetic particles, although they can also be other magnetic metal or metal oxides, whether in impure, alloy, or composite form, as long as they have a reactive surface and exhibit an ability to react to a magnetic field. Other materials that may be used individually or in combination with iron include, but are not limited to, cobalt, nickel, and silicon. Methods of making magnetite or metal oxide particles are disclosed in Vandenberghe et al., "Preparation and Magnetic Properties of Ultrafine Cobalt Ferrites," J. of Magnetism and Magnetic Materials, 15 through 18: 1117-18 (1980); E. Matijevic, "Mono Dispersed Metal (Hydrous) Oxide--A Fascinating Field of Colloidal Science," Acc. Chem. Res., 14:22-29 (1981), the disclosures which are incorporated herein by reference.

15 A magnetic bead suitable for application to the present invention includes a magnetic bead containing primary amine functional groups marketed under the trade name BIO-MAG by Advanced Magnetics, Inc. A preferred magnetic particle is nonporous yet still permits association with a probe moiety. Reactive sites not involved in the association of a probe moiety are preferably blocked to prevent nonspecific binding of other reagents, impurities, and cellular material. The magnetic particles preferably exist as substantially colloidal suspensions. Reagents and substrates and probe moieties associated to the surface of the particle extend directly into the solution surrounding the particle. Probe moieties react with dissolved reagents and substrates in solution with rates and yields characteristic of reactions in solution rather than rates associated with solid supported reactions. Further, with decreasing particle size the ratio of surface area to volume of the particles increases thereby permitting more functional groups and probes to be attached per unit weight of magnetic particles.

35 Beads having reactive amine functional groups can be reacted with polynucleotides to covalently affix the

462150 0808E280

polynucleotide to the bead. The beads are reacted with 10 percent glutaraldehyde in sodium phosphate buffer and subsequently reacted in a phosphate buffer with ethylenediamine adduct of the phosphorylated polynucleotide in a process which will be set forth in greater detail in the experimental protocol which follows.

Returning now to Step 2, the retrievable support with associated probe moieties is brought into contact with the clinical sample and, progressing through to Step 3, is brought into binding conditions. The probe moiety specific for the target of interest becomes bonded to the target strands present in the clinical sample. The retrievable support, dispersed throughout the sample and reagent medium, allows the probe moieties and target to hybridize as though they are free in a solution.

Hybridizations of probe to target can be accomplished in approximately 15 minutes. In contrast, hybridizations in which either the probe or target are immobilized on a support not having the capability to be dispersed in the medium may take as long as 3 to 48 hours.

Extraneous DNA, RNA, cellular debris, and impurities are not specifically bound to the support. However, as a practical manner, a small amount of extraneous DNA, RNA, cellular debris, and impurities are able to and do in fact nonspecifically bind to any entity placed within the reaction vessel including the retrievable support. Embodiments of the present invention facilitate the further purification of clinical samples to remove extraneous DNA, RNA, cellular debris, and further impurities from target polynucleotides.

Step 4 of Figure 1 depicts the separation of the support of the clinical sample and the suspension of the support into a second medium. The second medium thus includes the retrievable support with the associated probe bound to target polynucleotide strands. Also carried with the retrievable support is extraneous DNA, RNA,

cellular debris, and impurities nonspecifically bound to the support, but in a much lower concentration than what was initially found in the clinical sample. Those skilled in the art will recognize that some undesirable materials can be reduced by washing the support prior to suspension in the second medium.

The retrievable support with associated probe and target strands suspended in the second medium is subject to further denaturation as set forth in Step 5 thereby allowing the target to disassociate from the probe moieties of the retrievable support. The denaturation process may or may not release nonspecifically bound extraneous DNA, RNA, cellular debris, or impurities from the retrievable support. However, Step 5 of the present method allows the retrievable support to be removed from the second medium carrying with it much of the nonspecifically bound cellular debris, impurities, and extraneous DNA, and RNA initially carried over from the first clinical sample medium.

As set forth in Step 6, a new support can be introduced into the second medium under binding conditions to again capture target polynucleotide strands on probe moieties associated with the retrievable support. It will be recognized by those skilled in the art that the new support may actually include the original retrievable support after recycling steps to further purify and remove nonspecifically bound DNA, RNA, cellular debris, and impurities. Thus, the only impurities present in the second medium include DNA, RNA, cellular debris, and impurities previously nonspecifically bound to the support which has subsequently been released from the first support and dissolved or suspended in the second medium.

However, such impurities can be further removed from the target polynucleotides by removing the second retrievable support from the second medium and again repeating the cycle of introducing the retrievable support into a further medium, denaturation, and removal of

the old support. Those skilled in the art will recognize that the magnetic beads described in the present invention are susceptible of being raised out of a solution or being held in place as a solution is removed or added to a containment vessel.

The ability of the magnetic beads to participate in the reactions which mimic "insolution kinetics" strands allow the completion of a cycle of denaturation and binding to the target to be accomplished in three to fifteen minutes.

After sufficient purification and concentration, the target can be detected by luminescent or radioactive methods known in the art as indicated in Step 8. Purification of the medium containing the target allows the detection of nonisotopic label moieties without cellular debris and impurities.

Turning now to Figure 2, which features a multiple probe method, a further embodiment to the present assay method is illustrated beginning with a clinical sample containing polynucleotide target which is processed in accordance with the clinical sample of the previous figure with the introduction of solubilizing agents and reagent. The reagent of the assay method depicted in Figure 2 includes a first polynucleotide probe strand (P_1) and a second polynucleotide probe strand (P_2) capable of forming a complex with the target in which both probes (P_1 and P_2) are bound to the target. The first probe (P_1) is capable of associating with a retrievable support (S_1) under binding conditions. The second probe has at least one label moiety capable of detection. The label moiety is illustrated in the drawings with an asterisk or a star. Following the introduction of solubilizing agents and reagent under denaturation conditions, the solution containing the clinical sample potentially includes target polynucleotides and reagent in the form of the first and second probes, plus cellular debris, solubilizing agents, impurities, and

002234000 050094
46050 0805280

25

30

35

25

hybridization of the probes to target insolution, a solid nonbead or nonparticulate membrane or filter support can be incorporated into the reaction vessel. The solution of reagent and sample can be passed through the support to affect target capture. The support (S_1) is illustrated in Figure 2 as a retrievable support.

In solution with the target-probe support complex are unbound first and second probe moieties, unbound target solubilizing agents, impurities, and cellular debris. The unbound second probe (P_2) which has label moieties constitutes noise, producing a signal which mimics the presence of target. A small amount of extraneous cellular debris, solubilizing agents, impurities, and probes may also become nonspecifically bound to the retrievable support.

In Step 4, the support (S_1) is separated from the clinical sample medium. If a retrievable support is used, separation can be accomplished either by immobilizing the retrievable support within a reaction vessel or by withdrawing the retrievable support from the sample medium directly. Those skilled in the art will recognize that the immobilized support can be washed to reduce undesirable material.

Turning now to Step 5, the target-probe support complex is substantially free of extraneous RNA, DNA, solubilizing agents, impurities, and cellular material and can be monitored for the presence of the label moieties indicative of the presence of the target molecule. However, a small amount of extraneous DNA, RNA, solubilizing agents, impurities, and cellular materials may still be nonspecifically bound to the support (S_1). Moreover, unbound, in the sense that it is not associated with target, second probe (P_2) may also be nonspecifically bound to the support (S_1) and can affect signals from nonisotopic label moieties. The presence of unbound second probe moiety (P_2) having label moieties is a significant cause of background noise thereby reducing the

accuracy of the assay procedure.

Thus, as an alternative Step 5, the first support (S_1) may be suspended into a second medium where the support (S_1) is separated from the target-probe complex by denaturation.

Following denaturation, in Step 6, the first support (S_1) is removed from the second medium and replaced with a second support (S_2). The second support (S_2) includes an antiligand moiety (A_1) capable of binding to the ligand moiety (L_1) of the first probe.

Moving to Step 7, under binding conditions, the target-probe complex reassociates with the second support (S_2). The removal of the first support (S_1) removes extraneous material, debris, and probes nonspecifically bound to the first support (S_1) from the assay medium.

As illustrated in Step 8, the medium containing the target-probe complex can be monitored for the presence of the labels. However, further purification of the assay medium can be performed to further reduce the presence of background and extraneous materials which may have been carried from the sample medium nonspecifically bound to the first retrievable support (S_1) and subsequently dissolved or disassociated from the first support (S_1) into the second medium.

Thus, the second retrievable support (S_2) may be brought into contact with a third medium, the medium brought into conditions to release the target-probe complex from the support, and the support removed to complete a further cycle. The number of cycles will be a matter of choice depending on the type of sample, type of label moieties, and the sensitivity of the detection equipment. Different types of supports may be used at different times. Thus, a retrievable support can be used to gather or concentrate the target-probe complexes from sample mediums or solutions initially to avoid problems of clogging typical of membranes or filters. The second or third supports preferably includes a membrane or

00230080-050794
+05050-08082280

filter with antiligand moieties (A_1) which bind to the ligand moiety (L_1) of the first probe (P_1). Membrane or filter supports can simplify process steps allowing flow-through recovery of target-probe complexes.

5 A further embodiment of the present invention is particularly well suited for reducing background noise. Referring now to Figure 3, a modification of the previous assay procedure illustrated in Figure 2 is described. In Figure 3, a target polynucleotide has formed a complex
10 with a first and second probe moiety (P_1 and P_2) similar to the probe moieties described in Figure 2. However, the second probe includes a second ligand (L_2). The second ligand (L_2) may include, by way of example, a single terminal ribonucleotide which complexes with a
15 borate antiligand, an alternating copolymer which binds with a complementary copolymer, a biotin ligand which binds to an avidin antiligand, or as illustrated, homopolymer ligand (L_2), and a complementary homopolymer anti-
ligand (A_2).

20 Turning now to Step 1, a background support capable of selectively binding to the second probe (P_2), only when it is not bound to a target, is brought into contact with the medium containing the target-probe complex. The medium further includes free, disassociated first and
25 second probes (P_1 and P_2). The labeled second probe (P_2), which contributes to the background noise, is specifically bound to the background support (B_1) by a vast molar excess of antiligand moieties (A_2) associated with the background support (B_1). Following binding of the
30 unbound labeled probe (P_2) to the background support (B_1), the background support (B_1) is removed from the medium as illustrated in Step 2. The medium containing the target-probe complex can be monitored for the presence of the label contained upon the second probe (P_2)
35 with a reduction in background noise. Alternatively, the medium containing the target-probe complex can be subjected to further processing.

5

10

25

30

35

5

10

15

20

30

35

form in Figure 7. The apparatus includes the following major elements: at least one containment vessel, means for controlling the association of a probe with a target molecule and a retrievable support, means for separating the retrievable support from a sample solution, and means for releasing the target molecule from the retrievable support. These major elements may take various forms and are described more fully below.

The apparatus will be described below for illustration purposes as applying the methods described in Figures 2 and 3 relative to a target molecule which includes a polynucleotide. Thus, at Station 1, a clinical sample is placed within the containment vessel with solubilizing agents such as chaotropic salts, enzymes, and surfactants in order to dissolve cellular material and release nucleic acids. The containment vessel may include agitation elements to facilitate the break up of cells. The containment vessel may include any type of vessel, tube, cuvette suitable for containing the sample.

In an instrument designed for automated analysis, the apparatus set forth in Figure 7 will preferably include means for receiving a plurality of containment vessels. For illustration purposes, the containment vessels containing the sample are analyzed sequentially. Thus, containment vessels are conveyed to a first station and then to subsequent stations where various steps of the assay method are performed.

The various stations are linked by conveying means. Conveying means may include a rotatable turntable, conveying belt, or the like. As applied in a clinical hospital setting, conveying means may include manual movement. Thus, hospital staff may obtain a tissue sample from a patient and place the sample in the containment vessel. Sample processing, including the breakup of the tissue sample and initial mixing of solubilizing agents and reagents would be initiated at bedside and continued as the containment vessel traveled to a subsequent sta-

00232080-050394

tion for further processing. References to stations are for illustration purposes. Those skilled in the art will recognize that certain stations or steps may be combined or reversed.

5 Returning now to the first station, sample and solubilizing agents are placed within a containment vessel in which an agitation element thoroughly mixes the sample and solubilizing agents, releasing nucleic acids from cellular materials. Conveying means carry the contain-
10 ment vessel to Station 2 where the containment vessel receives reagent.

15 The reagent includes a first polynucleotide probe and a second polynucleotide probe. The first and second probes are capable of forming a complex with the target polynucleotides in which both probes are bound to mutu-
20 ally exclusive portions of the target. The first probe is also capable of binding to a retrievable support under binding conditions. The second polynucleotide probe includes a label moiety capable of detection. The
25 reagent and sample nucleic acid are denatured by a heating element and conveyed to Station 3.

30 At Station 3, the containment vessel receives a first support depicted by open circles. The first support is homogeneously dispersed within the sample
35 medium by suitable means including an agitation element. Examples of suitable supports include, without limita-
40 tion, polystyrene beads, magnetic beads and other particulate or filamentous substances. As illustrated, the first support includes a magnetic bead having polynucleo-
45 tide antiligands of deoxythymidine (dT). The first probe includes a tail portion of deoxyadenosine (dA) capable of binding to the first support during binding or hybridiza-
50 tion conditions.

55 Moving to Station 4 hybridization conditions are imposed upon the sample medium by cooling by a cooling
60 element. However, those skilled in the art will recognize that means to alter salt concentrations can be

44-38867-10000

readily substituted for thermal controls. Thus, the target polynucleotide forms a complex with the first and second probes. Further, the homopolymer deoxyadenosine (dA) tail portion of the first probe hybridizes to the deoxythymidine (dT) homopolymer of the retrievable support.

From Station 4, the containment vessel is moved to Station 5 where the retrievable support is immobilized on the wall of the containment vessel by activating a magnetic element. If polystyrene beads were substituted for magnetic beads, the polystyrene bead would be immobilized by filtering or density differences. The sample medium is disposed of carrying with it most of the extraneous DNA, RNA, solubilizing agents, cellular material, and impurities. The immobilized retrievable support is washed to further remove extraneous DNA, RNA, solubilizing agents, cellular materials, and impurities.

Further, although it is illustrated that the retrievable support is immobilized on the wall of the reaction vessel, it is also possible to remove the retrievable support from the reaction vessel by a magnetic element and dispose of the first reaction vessel containing with it extraneous DNA, RNA, solubilizing agents, and cellular material which may be nonspecifically bound to the reaction vessel walls.

The retrievable support is placed in a second medium, either the same containment vessel or a new containment vessel. The containment vessel, containing the retrievable support in a second medium is carried to Station 6.

At Station 6, the second medium is brought to denaturation conditions by suitable means including a heating element. The denaturation process releases the target-first and second-probe complex from the (dT) homopolymer of the retrievable support. The first support, potentially carrying extraneous DNA, RNA, impurities, and cellular material, is removed from the second medium. If

460050-0808F280

desired, amplification steps may be applied to the target, now substantially free of impurities, debris, and non-target polynucleotides. Amplification steps may include the generation of an amplification product with enzymes such as, by way of example, DNA polymerase, RNA polymerase, transcriptases, or Q β replicase. In the event the amplification product is not the target molecule, the second probe is directed to the amplification product as well as a third capture probe which takes the place of the first probe. A background support is then brought into contact with the second medium and passed to Station 7.

At Station 7 a cooling element brings the second medium to hybridization temperatures. The background support includes a second antiligand capable of specifically binding to a ligand carried upon the second probe. For example, without limitation, a terminal nucleotide of the second probe can be synthesized to be a ribo derivative which specifically binds to borate moiety carried upon the second support. The second probe bound to the target as part of a probe target complex will not bind to the borate carried upon the third support due to steric hindrances. However, unbound second probe will specifically bind to the borate support. Alternatively, the second probe may include a homopolymer such as deoxycytosine (dC) which binds to a deoxyguanine (dG) homopolymer linker on a second support. The length of the homopolymers are designed such that complexes of the target-first and second probes with the second support are not stable; however, complexes of the second probe alone with the second support are stable within reaction parameters. Indeed, background capture binding of background support to unbound second probe can be irreversible.

Next, the containment vessel containing the second medium and the background support is conveyed to Station 8 where the background support having second probe strands unbound to the target-probe complex is separated

from the second medium. Separation of the background support removes nonspecific background noise from the medium.

5 As illustrated, background capture is effected upon beads. However, those skilled in the art will recognize that the initial purification of the target-first and second probe complex from the clinical sample, removes all or most solid debris allowing background capture on filter or membrane supports through which the second
10 medium can be flushed.

From Station 8, the purified medium containing the target-probe complex with reduced background is conveyed to Station 9. At Station 9, a third support, depicted as a membrane or filter, is brought into contact with the
15 second medium which is brought to hybridization temperatures by a heating element. The third support includes first antiligand moieties which bind to the first ligand moieties of the first probe, or if an amplification product is generated in previous steps, to a first ligand
20 moiety of a third probe directed to the amplification product. Thus, if the first ligand moiety of the first probe is of a homopolymer of deoxyadenosine (dA), the third support may include homopolymer of deoxythymidine (dT). As illustrated, the third support includes filters
25 or membranes through which the second medium can be flushed; however, beads or particles may also be used. The third support serves to further concentrate the target-first and second probe complex and permits further reduction of background and interfering materials which
30 do not specifically bind to the third support. Moving to Station 10, the third support concentrates the target-first and second probe complex allowing detection of label moieties carried upon the second probe.

35 The present invention is further described in the following typical procedures and experimental examples which exemplify features of the preferred embodiment.

440000 150000 440000 150000

35

I. Procedures

A. Materials

5 All reagents were of analytical grade or better. Magnetic beads marketed under the trademark BIO-MAG containing functional amino groups were obtained from Advanced Magnetics, Inc. of Cambridge, MA.

10 In the present example, all labeled nucleotides were obtained from New England Nuclear. The enzyme terminal deoxynucleotidyl transferase (TDT) was obtained from Life Sciences, Inc., St. Petersburg, Florida. The oligonucleotide pdT₁₀ was obtained from Pharmacia PL Biochemicals.

15

B. Synthesis of Probes

20 The following sets forth typical protocols and methods. Referring now to Figure 8, two probes were constructed to the sense strand of the enterotoxin gene elt A1 of Escherichia coli, in accordance with the constructional map, Figure 8, of Spicer, E. K. and J. A. Noble, 1982, J. of Biological Chem. 257, 55716-55751.

25 One set of probes was synthesized beginning at position 483 of the gene sequence and extending onward 30 nucleotides in length, hereinafter referred to as the A483 probe. A second probe was synthesized beginning at position 532 in the gene sequence and extending 30 nucleotides in length, hereinafter referred to as the A532 probe. A third probe was synthesized beginning at position 726 in the gene sequence and extending 39 nucleotides in length, hereinafter referred to as the A726 probe. The specific base sequences (5' to 3') are set forth in Table 1 below:

35

Table I

Probe	Sequence
A483	AGA CCG GTA TTA CAG AAA TCT GAA TAT AGC
A532	AGA TTA GCA GGT TTC CCA CCG GAT CAC CAA
A726	GTC AGA GGT TGA CAT ATA TAA CAG AAT TCG GGG GGG GGG

The probes were synthesized by methods available in the art. The numbering system is adapted from the 768 nucleotide sequence available through Intelligenetics sequence bank ECO ELT A1.

Of the ten G residues at the 3 prime end of probe A726, three guanine bases towards the 5' end are capable of binding to three complementary cytosine bases of the tox gene. Stretches of three cytosines are common in DNA. The ten guanine bases form a ligand capable of binding to a poly C antiligand carried upon a support such as oligo dC-cellulose. However, seven guanine bases will not form a stable association with a support at 37°C, particularly if the probe is bound to target due to steric hindrance and the size of the target-probe complex. Probe A726 was modified by the random addition of approximately three residues of ³²P-dC and ³²P-dG to its 3' end with terminal transferase.

Those skilled in the art will recognize that other probes can be readily synthesized to other target molecules.

C. Preparation

The target in Example Nos. 1, 2 and 3 is the enterotoxin gene elt A1. The enterotoxin gene elt A1 is carried as part of the plasmid EWD-299 obtained from Stanford University.

In Example No. 1, enterotoxigenic bacteria were grown to log-phase in Luria broth. The enterotoxigenic bacteria were lysed and the plasmid EWD-299 isolated.

5 The plasmid EWD-299 was further digested with the restriction enzymes Xba 1 and HIND III. A fragment of 475 base length was used as a target and purified by electro-elution from a 1 percent agarose gel. In order to follow the efficiency of capture steps, the fragment was 5' end labeled with ^{32}P -ATP with the enzyme polynucleotide kinase following manufacturer's instructions.

10 In Example Nos. 2 and 3, the enterotoxigenic bacteria and wild type nonenterotoxigenic E. coli JM83 were separately grown to log phase. The wild type E. coli serves as a control. Separate extracts of enterotoxigenic bacteria and wild type bacteria were prepared by substantially solubilizing the cells in chaotropic solutions. Thus, the bacteria cultures, in Luria broth, 15 were added to solid guanidinium thiocyanate (GuSCN) to a concentration of 5M GuSCN, Tris-HCl to a concentration of 0.3M, and EDTA (pH7) to a concentration of 0.1M. The chaotropic-bacterial solutions were then heated to 100°C for five minutes and cooled. The resultant enterotoxi- 20 genic bacteria extract was serially diluted with wild type nonenterotoxigenic bacteria extract. The concentration of tox plasmids per cell and the cell number in the extracts were measured by conventional techniques. The original extracts solubilized in GuSCN contained approxi- 25 mately 10^9 enterotoxigenic E. coli per ml and 100 plasmids/cell.

D. Synthesis of Beads

30 Retrievable supports were prepared from magnetic beads. Other retrievable supports include particles, fibers, polystyrene beads or other items capable of physical separation from a medium. Magnetic beads were synthesized with an adduct of deoxythymidine of ten base length to allow the beads to associate with probes tailed 35 with deoxyadenosine in a readily reversible manner.

Thus, 100 ml of beads having amine functional groups such as BIO-MAG (M4100) beads were washed four times with

20 mM sodium phosphate (pH 6.7) in four 275 ml T-flasks. The beads were then washed with 1% glutaraldehyde in 20 mM sodium phosphate. Next, the beads were reacted in 100 ml of 10 percent glutaraldehyde in 20 mM sodium phosphate (pH 6.7) for three hours at room temperature. The beads were then washed extensively with 20 mM sodium phosphate (pH 6.7) and then washed once with 20 mM phosphate (pH 7.6).

Separately, a purified ethylene diamine (EDA) adduct of pdT_{10} (EDA-dT_{10}) was prepared in accordance with Chu, B.C.F., G. M. Wahl, and L. E. Orgel; Nucleic Acid Res. 11, 6513-6529 (1983) incorporated by reference herein. The concentration of EDA-dT_{10} was adjusted to 1 OD/ml in 20 mM phosphate (pH 7.6).

The EDA-dT_{10} was combined with the magnetic beads to allow the EDA-dT_{10} to react with the free aldehyde groups of the beads. The mixture of EDA-dT_{10} and beads was divided into a plurality of 50 ml polypropylene tubes. The tubes containing the reaction mixture and beads were placed in a tube rotator and agitated overnight at room temperature.

Next, the beads were washed five times to remove noncovalently bound EDA-dT_{10} with a wash solution of sterile 20 mM phosphate (pH 6.7) in large 275 ml T-flasks and diluted to 200 ml with the wash solution.

For storage, beads can be maintained for months in a buffer of 20 mM phosphate, to which is added sodium azide to 0.1% and SDS to 0.1%. Bead preparations are stored at 4°C protected from light.

The beads were then prehybridized to block nonspecific binding sites in a buffer, hereafter referred to as "prehybridization buffer", of 0.75 M sodium phosphate (pH 6.8), 0.5% sodium lauroyl sarcosine, 10 micrograms/ml E coli DNA, 0.5 milligram per milliliter mg/ml bovine serum albumin (BSA) (Nuclease-free) and 5 mM ethylenediaminetetraacetic acid (EDTA). Before applying the probes and beads to target capture procedures, two prehy-

145050 0308F280

bridizations of the beads were performed. The prehybridization procedure included placing the beads in ten volumes prehybridization buffer.

5 The first prehybridization procedure was performed with agitation at 60°C. The second prehybridization procedure was performed at room temperature with swirling. A 0.1 percent isoamyl alcohol solution was added to the solutions as a defoamant.

10 The binding capacity of dT₁₀-derivatized beads was measured by the following procedure. In separate vessels, dT₅₀ and dA₅₀ were 5' end labeled with ³²P-dT and ³²P-dA respectively to a specific activity (Sa) of about 10⁶ dpm/microgram. Next, the Sa was accurately measured for a known quantity of reacted dT₅₀ by trichloroacetic acid precipitation.

15 Next, 5 µg of ³²P-dA₅₀ and 5 µg of ³²P-dT₅₀, having substantially identical Sas of between 100,000-200,000 dpm/mg, were separately added to tubes containing prehybridization buffer and brought to a volume of 1 ml.

20 A known sample volume of prehybridized beads was placed into four tubes. Two of the four tubes each receive 0.5 ml of the ³²P-dA₅₀ mixture and the remaining two tubes receive 0.5 ml of the ³²P-dT₅₀ mixture. All four solutions are brought to hybridization conditions for five minutes. The beads are thereafter immobilized and washed. The activities of the solutions are then monitored. The total binding capacity, C, for a quantity of bead preparation measured in micrograms is set forth below:

30
$$C = V(A - T)/X$$

In the above equation X is the specific activity of ³²P-dT₅₀ in cpm/mg, V is the volume ratio of total volume to sample volume, A is the average activity of the beads suspended in ³²P-dA solutions in cpm, and T is the average activity of the beads suspended in ³²P-dT solutions in cpm.

08238080-050344

40

Those skilled in the art will recognize that other beads, particles, filaments, and the like can be formulated with other nucleotide combinations or homopolymers. For example, polyA-derivized beads were produced by substituting (for the purified EDA adduct of dT₁₀) a solution containing 100 mg poly A (mw>100,000) in 50 ml of 20 mM sodium phosphate (pH7.6).

E. Target Capture Procedures

Bead preparations were used to capture target polynucleotides. The following sets forth a typical experimental target capture protocol demonstrating retrievable supports and reversible captures for purposes of illustration, without limitation, the procedure will be discussed using a first probe A483 and a second probe A532. The first probe, A483, was randomly 3' end labeled with ³²P-dCTP and ³²P-dGTP to a specific radioactivity of about 10¹⁰ dpm/mg. The second probe, A532, was tailed with about 70 unlabeled dA residues by the enzyme terminal transferase.

First, 200 µg/ml of labeled probe A483 and 400 µg/ml of tailed probe A532 were mixed with varying amounts of a heat-denatured 475 mer Xba I-HIND III restriction fragment of the enterotoxin gene at 65°C for 15 minutes in 1.4M sodium chloride.

Next, target capture was initiated by contacting the medium containing the target and probe moieties with an aliquot of dT₁₀-magnetic beads having 3 micrograms/ml of dA₅₀ binding capacity following prehybridization procedures to reduce nonspecific binding to the magnetic bead. The magnetic bead and the probe-target complex was incubated at room temperature in 0.1 ml prehybridization buffer in 5 ml polypropylene tubes for two to five minutes.

The tubes were placed into a Corning tube magnetic separator. The Corning tube magnetic separator upon activation imposes a magnetic field through the polyprop-

44-059-0786300

44

pylene tubes which immobilizes the magnetic beads on the inner walls of the tubes. During the time that the magnetic beads are immobilized on the side walls of the polypropylene tubes, the original medium was removed and discarded.

While immobilized, the beads were washed three times with 0.6 ml of prehybridization buffer containing isoamyl alcohol as a defoamant. Following the addition of the prehybridization buffer, the beads were resuspended by removing the tubes from the magnetic field and by subjecting the medium to vigorous vortexing.

Next, the magnetic field was reapplied to immobilize the beads allowing the prehybridization buffer to be removed and discarded. The cycle of adding the prehybridization buffer, resuspending the beads, immobilizing the beads, and discarding the prehybridization buffer was repeated twice. Target-probe complexes held on the beads are available for further processing including additional steps of detection, background capture or further cycles of target capture.

A preferred target capture procedure includes release of the target-probe complex and recapture on a second support. Preferably the support is chemically distinct from the first support.

Release of the target-probe complex is effected in the following typical protocol. After the removal of the last prehybridization buffer, prehybridization buffer was added to the tube containing the beads. The beads were incubated with agitation at 60°C for one-two minutes to release the probe-target complexes from the bead. The magnetic separator was again activated with the temperature at 60°C and the eluate, containing free target-probe complexes, is removed from the tube. The eluate can be recaptured on additional retrievable supports or subjected to final capture on conventional supports. It will be recognized by those skilled in the art that the capture and release of the target probe complex from ret-

retrievable supports such as the magnetic beads of the present example can be repeated as often as desired to reduce hybridization backgrounds.

Final capture of the probe-target complex was typically performed on nitrocellulose filters or nylon membranes containing nonspecifically bound or covalently bound dT-3000. Thus, the target-probe complexes carried upon the magnetic beads were released from the magnetic beads by heating the beads to 60°C in prehybridization buffer for two minutes. The beads were immobilized and the eluate removed and passed through a 0.2 micron acro-disc (Gelman) to remove magnetic fines. The nitrocellulose filter containing dT-3000 selected, bound, and captured the dA tail on the unlabeled probes.

The use of a chemically different solid support for the final capture of the target-probe complex avoids binding background molecules which may have a high affinity for previously used supports. By way of illustration, it is possible for lower level contaminants with a natural high affinity for a particular support to repeatedly bind and elute with a support along with probe-target complexes. Such low level contaminants cannot be diluted out by repeated use of a retrievable support of the same composition as completely as by exposing them to supports of very different compositions. Low level contaminants can also be lowered by utilizing chemically distinct means to release the target-probe complexes from supports and recapture.

F. Background Capture Procedures

Background capture procedures permit the selective reduction of background noise permitting the detection of true signal indicative of the presence of target. Background capture can be applied in a single probe system or in systems using more than two probes. For example, in background capture procedures featuring a single probe,

46E050-0809E730

the probe includes a label moiety and a ligand. The probe is capable of binding to a target and the ligand is capable of forming a stable bond to a support only when the probe is unbound to target.

5 Similarly, by way of example, background capture procedures featuring multiple probes in conjunction with target capture include two probes. A first target capture probe, having an unlabeled ligand capable of binding to a first support is used to capture the target and a
10 second background capture probe, having a label moiety capable of detection includes a second ligand capable of binding to a second background support. Background capture is a valuable supplement to target capture for enhancing the signal to noise data of an assay.

15 The following sets forth a typical background capture protocol using a first target capture probe A532 and a second background capture probe A726 and a target enterotoxin gene elt A1. Those skilled in the art will recognize that the probes used for demonstration purposes
20 are merely a matter of choice. Other probes could be used also.

45C050" 0808E280
25 The probe A532 was tailed with approximately 100 dA residues capable of reversibly binding to dT₁₀ covalently linked magnetic beads for initial target capture and dT₃₀₀₀ nonspecifically bound to nitrocellulose for a final target capture. The probe A726 was end labeled with the random addition of approximately three residues of ³²P-dC and ³²P-dG to the 3' end with terminal transferase. The probe A726 is capable of binding to dC-cellulose when the probe is not hybridized to target.
30

35 A solution containing the target-first and second-probe-complex and potentially containing unbound second probe is mixed with dC-cellulose and the temperature of the mixture maintained at 37°C. The temperature, 37°C, is higher than the dissociation temperature of dG₇ with oligo dC, preventing binding of the target-first and second-probe-complex to the dC-cellulose. The temperature

44

is also lower than the dissociation temperature of dG₁₀ with oligo dC to promote binding of unbound second probe having a dG tail to the dC-cellulose. Additionally, the target-first and second probe complex is sterically hindered to a greater degree in its approach to the dC-cellulose support than unbound second probe. The dC-cellulose containing the second probe A726 is removed by centrifugation, however, those skilled in the art will appreciate that other methods such as filtration may be used as well. The remaining eluate contains target-first and second probe complexes and a reduced concentration of unbound labeled second probe A726.

G. Examples

Individuals skilled in the art will recognize that the typical protocols for retrievable support preparation, probe preparation, target capture and background capture are capable of modification to suit special needs and purposes. The following examples incorporate the typical procedures outlined above unless otherwise noted.

Example 1

Target Capture and Assay Using Magnetic Bead

A target capture assay was performed with two probes and a magnetic bead retrievable support. The target included the Xba I-Hind III fragment of the enterotoxigenic gene elt A1. A first probe included an A532 thirtimer oligonucleotide probe which was tailed with 130 unlabeled dA residues capable of binding to the dT₁₀ residues of the magnetic beads support. A second probe included an A483 thirtimer oligonucleotide probe capable of binding to the same target 20 nucleotides downstream from the site of hybridization of the first probe. The second probe was labeled by tailing the thirtimer oligonucleotide with ³²P-dCTP and ³²P-dGTP to a specific radioactivity of 10¹⁰ DPM/microgram.

465050" 0808E780

465

The tailed first probe and the labeled second probe were incubated at 65°C for 15 minutes in 1.4 M sodium chloride with various quantities of heat denatured 475mer restriction fragments of the tox gene. As a nonspecific binding background control, the tailed first probe and labeled second probe were incubated in identical solutions in the absence of any target. As specific binding controls, two additional reaction mixtures were formed. One reaction mixture included the tailed first probe and the unlabeled second probe incubated with four micrograms of denatured E. coli DNA, and a second reaction mixture of the tailed first probe and the labeled second probe incubated in ten micrograms of denatured human DNA in identical reaction mixtures without any target DNA.

After a 15 minute hybridization period, the samples were incubated for five minutes with dT-derivatized magnetic beads in 0.7 milliliters of 0.75 molar phosphate buffer (pH 6.8). The beads were magnetically immobilized and washed extensively as described previously. The target-probe complex was eluted from the beads at 60°C in 0.6 milliliters of 0.20 molar phosphate buffer (pH 6.8). The first set of beads was separated from the eluate and the target probe complex. A second group of magnetic beads was added to the eluate and brought to binding conditions to capture the target and probe complex again. The second set of beads was washed and the target again eluted from the beads and the beads separated from the eluate.

A third set of beads was added to the eluate containing the target-probe complex and placed under binding conditions to allow the beads to once again capture the target-probe complex. The beads were then washed extensively and the target eluted from the beads as previously described. The beads were then separated from the eluate and the eluate passed through dT₃₀₀₀-nylon into two millimeter square slots, capturing the target-probe complex.

46

46E.050" 7803F.280

5 The dT₃₀₀₀ nylon membrane was prepared in which 2 µg dT₃₀₀₀ was covalently bound to nylon using a hybri-slot apparatus (Betlesda Research Laboratory). Briefly, dT₃₀₀₀ (Life Sciences) was dotted directly onto a nylon membrane such as Gene-ScreenTM (New England Nuclear) in a salt-free Tris buffer. The membrane was dried at room temperature for 10 minutes, and then dried under an infrared lamp for an additional 10 minutes before cooling back to room temperature for another 10 minutes. The filter apparatus containing the nylon membrane was inverted on a uv-transilluminator (Fotodyne) and exposed to uv light for two minutes at 40uW/cm² to cross-link the dT₃₀₀₀ to the filter.

15 The dT₃₀₀₀ membrane was prehybridized by sequentially passing the following solutions through the membrane:

- (1) 1% SDS;
- (2) 0.5 mg/ml BSA in 0.5% SDS; and, finally,
- (3) prehybridization buffer

20 The dT₃₀₀₀-nylon potentially containing the target-probe complex was washed with 0.2 molar sodium phosphate and 5 millimolar EDTA. The nylon support was monitored overnight by audioradiography for the presence of the ³²P label moieties of the second probe. Following audioradiography, the bands were cut out of the filter and counted in base scintillation fluid. The counts were 2100 and 1400 counts per minute in the solution containing three femtomoles (10⁻¹⁵ moles) of a restriction fragment containing the tox gene. Samples containing 30 attomoles (10⁻¹⁸ moles) of the restriction fragment containing the tox gene produced a count of 62 counts per minute.

35 A third sample containing no DNA produced seven counts per minute. A fourth sample containing ten micrograms of heat denatured human DNA produced 0 counts per minute. A fifth solution containing 4 micrograms of heat denatured E. coli DNA produced 7 counts per minute. The absolute sensitivity of the protocol was estimated to be

10⁻¹⁸ mole of tox gene. The overall efficiency of the recovery of labeled target-probe complex was estimated to be 1 to 2 percent of the input. The assay demonstrated good specificity. There is no more labeled probe in the samples containing human DNA or E. coli DNA than in the sample containing no DNA at all. Repetition of the experimental protocol has produced overall efficiency of capture of the target of almost 5 percent. The procedures reduced background from an initial level of 10¹¹ molecules of the labeled unhybridized probes to about 10⁴ moles. The reduction and background represents a 7 log improvement which more than adequately compensates for the reduction and efficiency of capture.

Example 2

The present example features target capture with background capture. Target and background capture was effected using an unlabeled first target capture probe, A532 as described in target capture, and a second labeled background capture probe A726.

First, 160 ng/ml dA-tailed A532 and 40 ng/ml ³²P-labeled probe A726 were combined to form a probe mix. The probe mix was added to 5 µl of bacterial extract containing various amounts of enterotoxigenic gene. The extract-probe mix was incubated at 22°C for 15 minutes.

After a fifteen minute hybridization period, the samples were diluted with ten volumes of prehybridization buffer incubated for five minutes with dT-derived magnetic beads in 0.7 ml of .75M phosphate buffer (pH 6.8) to effect target capture. The beads were magnetically immobilized and washed extensively. The target-first and second probe complex was eluted from the first support as previously described and the first solid support removed.

Next, the eluate containing the target-first and second-probe-complex and potentially containing unbound second probe was mixed with dC-cellulose and the temperature of the mixture maintained at 37°C. The temperature

44-557-0803E280

48

5
10

15

20

T490

Second Experiment

49

5

10

20

25

35

50

spiked biological media. The spiked biological media resembles samples which would be obtained clinically in a medical setting.

Cell extracts of enterotoxigenic E. coli and wild type E. coli were prepared as previously described. To measure the sensitivity of the detection of tox genes in an environment analogous to a clinical setting, extract containing toxigenic bacteria was diluted with the extract containing the wild type E. coli as previously described.

The following materials were obtained from anonymous donors: human stool sample, cow's milk, human saliva, human phlegm, human whole blood, human serum, human urine and human semen. Clinical-type samples were solubilized; over a time period of ten minutes. The stool sample, due to its solid nature, was solubilized in a solution of 5 M GuSCN, 0.3 M Tris-HCl (pH 7.4), 0.1 M EDTA (pH 7), 1% betamercaptoethanol. Following solubilization, aliquots of the sample were made and each aliquot was spiked with a known quantity of either toxigenic E. coli or wild type E. coli. The mixture was then passed through a crude filtration (Biorad Econocolumn) and heated to 100°C for five minutes.

The remainder of the samples were more liquid in nature and were handled differently than stool. Liquid samples were added to solid GuSCN to make the final concentration 5M. The solid GuSCN also contained sufficient Tris-HCl, EDTA, and betamercaptoethanol to make the final concentrations the same as in the stool example. Next, aliquots of the samples were made and each aliquot was spiked with a known amount of toxigenic E. coli or wild type E. coli. The mixture was passed through a crude filtration (Biorad Econocolumn) and heated to 100°C for five minutes.

The preparation of probes in Example 3 differs from previous examples. A first capture probe was generated with the plasmid pBR322. The plasmid was restricted with

46C.D50" 0202E230

51

S

15

20

35

1996-1997		1997-1998		1998-1999		1999-2000		2000-2001		2001-2002		2002-2003		2003-2004		2004-2005		2005-2006		2006-2007		2007-2008		2008-2009		2009-2010		2010-2011		2011-2012		2012-2013		2013-2014		2014-2015		2015-2016		2016-2017		2017-2018		2018-2019		2019-2020		2020-2021		2021-2022		2022-2023		2023-2024		2024-2025		2025-2026		2026-2027		2027-2028		2028-2029		2029-2030		2030-2031		2031-2032		2032-2033		2033-2034		2034-2035		2035-2036		2036-2037		2037-2038		2038-2039		2039-2040		2040-2041		2041-2042		2042-2043		2043-2044		2044-2045		2045-2046		2046-2047		2047-2048		2048-2049		2049-2050		2050-2051		2051-2052		2052-2053		2053-2054		2054-2055		2055-2056		2056-2057		2057-2058		2058-2059		2059-2060		2060-2061		2061-2062		2062-2063		2063-2064		2064-2065		2065-2066		2066-2067		2067-2068		2068-2069		2069-2070		2070-2071		2071-2072		2072-2073		2073-2074		2074-2075		2075-2076		2076-2077		2077-2078		2078-2079		2079-2080		2080-2081		2081-2082		2082-2083		2083-2084		2084-2085		2085-2086		2086-2087		2087-2088		2088-2089		2089-2090		2090-2091		2091-2092		2092-2093		2093-2094		2094-2095		2095-2096		2096-2097		2097-2098		2098-2099		2099-2100		2100-2101		2101-2102		2102-2103		2103-2104		2104-2105		2105-2106		2106-2107		2107-2108		2108-2109		2109-2110		2110-2111		2111-2112		2112-2113		2113-2114		2114-2115		2115-2116		2116-2117		2117-2118		2118-2119		2119-2120		2120-2121		2121-2122		2122-2123		2123-2124		2124-2125		2125-2126		2126-2127		2127-2128		2128-2129		2129-2130		2130-2131		2131-2132		2132-2133		2133-2134		2134-2135		2135-2136		2136-2137		2137-2138		2138-2139		2139-2140		2140-2141		2141-2142		2142-2143		2143-2144		2144-2145		2145-2146		2146-2147		2147-2148		2148-2149		2149-2150		2150-2151		2151-2152		2152-2153		2153-2154		2154-2155		2155-2156		2156-2157		2157-2158		2158-2159		2159-2160		2160-2161		2161-2162		2162-2163		2163-2164		2164-2165		2165-2166		2166-2167		2167-2168		2168-2169		2169-2170		2170-2171		2171-2172		2172-2173		2173-2174		2174-2175		2175-2176		2176-2177		2177-2178		2178-2179		2179-2180		2180-2181		2181-2182		2182-2183		2183-2184		2184-2185		2185-2186		2186-2187		2187-2188		2188-2189		2189-2190		2190-2191		2191-2192		2192-2193		2193-2194		2194-2195		2195-2196		2196-2197		2197-2198		2198-2199		2199-2200		2200-2201		2201-2202		2202-2203		2203-2204		2204-2205		2205-2206		2206-2207		2207-2208		2208-2209		2209-2210		2210-2211		2211-2212		2212-2213		2213-2214		2214-2215		2215-2216		2216-2217		2217-2218		2218-2219		2219-2220		2220-2221		2221-2222		2222-2223	
-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--



cally immobilizing the beads. The beads were washed twice during a fifteen minute time interval to remove impurities in the biological specimen and unhybridized biotin labeled second probe.

5 Next, in a time period of approximately one minute, the first and second probe-target complex was eluted from the magnetic beads at 65°C in blocking buffer. The eluate and the first beads were separated.

10 In a time period of approximately seven minutes, the first and second probe-target complex was releasibly bound to a second set of beads and again released. A second set of dT₁₀ derivized beads were then added to the eluate and hybridization conditions maintained for approximately one minute at 22°C. The beads were then
15 washed and resuspended in blocking buffer. The bead blocking buffer mixture was then brought to 65°C to release the first and second probe-target complex.

20 Over a time period of five minutes, final capture of the first and second probe-target complex on nitrocellulose was effected. The eluate from the second beads was filtered through a Gelman acrodisc (0.2 micron). The eluate containing the first and second probe-target complex was then passed through a dT₃₀₀₀ nitrocellulose filter (prehybridized with blocking buffer) at 22°C.

25 In a time period of approximately thirty minutes the filter was further processed to detect the biotin labels of the second probe. Buffer compositions used in detection are identified below in Table 3.

30

35

08238080 050394
+6E750 0808C280

Table 3
Detection Buffers

T940X

Buffer Number	Composition
1	1 M NaCl, 0.1 M Tris-HCl (pH 7.4), 5 mM MgCl ₂ , 0.1% Tween-20
1a	No. 1 with 5 mg/ml BSA, 10 micrograms/ml <u>E. coli</u> DNA
2	No. 1 with 5% BSA, 0.5% Tween-20
3	0.1 M NaCl, 0.1 M Tris-HCl (pH 9.5), 50 mM MgCl ₂

15 First, the filter carrying the first and second
probe-target complex, was incubated for approximately
five minutes in detection buffer No. 2. Next, the filter
was incubated for five minutes in a 1:200 dilution of
streptavidin-alkaline phosphatase (Bethesda Research Labo-
ratories) in detection buffer No. 1a. Thereafter, the
20 filter was washed three times in one minute in detection
buffer No. 1 and then washed twice in one minute in
detection buffer No. 3.

25 Next, 5-Bromo-4-chloro-3-indolyl phosphate (BCIP)
and nitroblue tetrazolium (NBT) (Kierkegaard and Perry)
were diluted twelve times in detection buffer No. 3, and
filtered through a 0.2 micron acrodisc. The diluted BCIP
and NBT solution was added to the filter and color
allowed to develop for fifteen minutes at 37°C.

30 Next, the filter was incubated in 50 mM Tris-HCl
(pH 7.4) and 10 mM EDTA for one minute to stop the reac-
tion. Sensitivity was determined visually on the filter
or by densitometric scanning on a CS 930 (Shimadzu Scien-
tific).

35 The steps in the present method are outlined below
in Table 4.

54

Table 4
Elapsed Time

T 9507

<u>Step Number</u>	<u>Time Required (min.)</u>	<u>Cumulative Timed (min.)</u>
1. Dissolution of biological sample; denaturation of DNA	10	10
2. Add labeled and unlabeled probes; hybridize in solution at 57°C	15	25
3. Capture probe-target complex on magnetic beads	1	26
4. Wash magnetic beads to remove impurities in the biological specimen and hybridization backgrounds	15	41
5. Elute the probe-target complex	1	42
6. Repeat steps 3-5 on a second set of beads (except abbreviate the washes)	7	49
7. Bind the probe-target complex to dT ₃₀₀₀ -nitrocellulose	5	54
8. Incubate filter in blocking buffer	5	59
9. Bind streptavidin-alkaline phosphatase	5	64
10. Wash	5	69
11. Add dyes to detect enzyme	15	84
12. Quench reaction	1	85

30 Although Table 4 set forth an example wherein the
elapsed time is just over one hour, the procedure is
capable of modification and can be performed in shorter
times. Nonradioactive probe assays of comparable sensi-
35 tivity may require twelve hours to several days and
require extensive sample preparation.

The sensitivity of the present assay is set forth in
Table 5 below:

95

Table 5

Sensitivity Level

<u>Biological Specimen</u>	<u>Concentration in the Hybridization Mixture</u>	<u>Number of Bacteria</u>
bacterial extract alone		1500
human stool	2.5% (w/v)	2000
cow's milk	12.5% (v/v)	3000
human saliva	12.5% (v/v)	3000
human urine	12.5% (v/v)	9000
human semen	2.5% (v/v)	9000
human blood	12.5% (v/v)	9000
human serum	12.5% (v/v)	9000
human phlegm	12.5% (v/v)	9000

20 Further, the present procedures are capable of further modifications to improve sensitivities. For example, a combination of thermal elution and chemical elution in multiple captured release cycles produces a signal to noise ratio five times better than single forms of elution, either multiple thermal elutions alone or

25 multiple chemical elutions alone.

Applying the same releasing or elution procedure tends to release the same background from the support. However, applying different releasing conditions tends to retain background on the support that would otherwise be

30 eluted. It is unlikely that background will behave identically to target under two physically or chemically distinct conditions.

A typical chemical elution of target-probe complexes on magnetic beads includes bringing beads in contact with 3M GuSCN for one minute at room temperature. Examples of thermal elutions have been described previously.

152050-0202280

56

The ability to detect bacteria would also be improved by directing probes to ribosomal RNA sequences. Ribosomal RNA sequences present a thousand fold increase in target per cell as compared to genomic DNA and clinically significant plasmid DNA.

The sensitivity of the above DNA or RNA target capture methods can be enhanced by amplifying the captured nucleic acids. This can be achieved by non-specific replication using standard enzymes (polymerases and/or transcriptases). After replication, the amplified nucleic acid can be reacted as above with capture probe, reporter probe, and capture beads to purify and then detect the amplified sequences.

In addition, where amplification is employed following purification of the target nucleic acids as described above, the amplified nucleic acids can be detected according to other, conventional methods not employing the capture probe, reporter probe, and capture beads described above, i.e., detection can be carried out in solution or on a support as in standard detection techniques.

Amplification of the target nucleic acid sequences, because it follows purification of the target sequences, can employ non-specific enzymes or primers (i.e., enzymes or primers which are capable of causing the replication of virtually any nucleic acid sequence). Although any background, non-target, nucleic acids are replicated along with target, this is not a problem because most of the background nucleic acids have been removed in the course of the capture process. Thus no specially tailored primers are needed for each test, and the same standard amplification reagents can be used, regardless of the targets.

The following are examples of the method.

35

57

Example 4:

The following example illustrates the use of RNA polymerase to amplify target DNA captured by a method which is a variation of the capture method discussed above.

Referring to Fig. 4, target DNA of a sample is first reduced in size by shearing or by limited nuclease digestion, according to standard methods. A recA protein coated capture probe is then added to the digested target DNA (Proc. Natl. Acad. Sci. U.S.A. (1986) 83:9591) The recA protein coated probe contains a nucleic acid sequence (a) that is homologous to a first target (a¹) sequence of the target DNA, as well as a homopolymer sequence homologous to a nucleic acid sequence on a capture bead. This capture bead is then added to the mixture to isolate and purify the target nucleic acid, as described above.

The captured DNA is amplified by treatment of the mixture with E. coli RNA polymerase lacking sigma subunit, i.e., core enzyme; E. coli RNA polymerase is described by R. Burgess in RNA Polymerase, Cold Spring harbor press, pp. 69-100, and can be purchased from New England Biolabs, Beverly, MA. The sigma subunit is removed according to the procedure described in J. Biol. Chem. (1969) 244:2168 and Nature (1969) 221:43. Other phage or bacterial RNA polymerases that lack transcriptional specificity can also be used. Core enzyme is added together with nucleotide triphosphates and a low salt transcription buffer such as described in Eur. J. Biochem. (1976) 65:387 and Eur. J. Biochem (1977) 74, 1107.

A suitable nucleotide triphosphate/transcription buffer solution has the following composition:

0 to 50 mM NaCl or KCl
25 mM Tris HCl pH 7.9 buffer
10 mM MgCl₂
0.1 mM EDTA
0.1 mM dithiothreitol

46E050 0808E280

58

0.5 mg/ml BSA

0.15 mM UTP, GTP, CTP, ATP

The resulting non-specific transcription of the target DNA produces many RNA transcripts of the target DNA which are then captured using a capture probe containing a sequence (b^1) homologous to a sequence (b) of the RNA transcripts. A reporter probe containing a sequence (c^1) homologous to another sequence (c) of the RNA transcript is then used for detection.

10 Example 5

In this example both non-specific replication of target DNA and transcription of that DNA are used to amplify captured target DNA.

Referring to Fig. 5, denatured sample DNA is captured as described above and the enzyme DNA polymerase (for example, Klenow fragment; Dur. J. Biochem, (1974) 45:623 available from New England Biolabs), random oligohexamer primers (i.e., Hexamers prepared to contain randomly selected bases at each nucleotide position in the hexamer) and deoxynucleotide triphosphates are added in appropriate buffers to cause replication of target DNA to form additional double stranded DNA. Suitable oligohexamer primers are available under catalog No. 27-2166 from Pharmacia, Inc. Piscotaway, NJ. A suitable deoxynucleotide triphosphate/buffer solution has the following composition:

- 66 mM glycine-NaOH buffer, pH 9.2
- 6 mM $MgCl_2$
- 1 mM 2-mercaptoethonal
- 30 mM each d CTP, d GTP, d TTP, d ATP

Because the primers are random, some will, simple as a matter of statistics, bind to and cause replication of sample sequences, no matter what those sequences are. (Alternatively, the double stranded DNA can be formed by synthesis starting from capture probe a.) RNA polymerase lacking sigma subunit is then added along with nucleotide triphosphates and low salt transcription buffer. Tran-

16E050" 0808E230

5 Example 6

Referring to Fig. 6, sample DNA is denatured, reduced in size and captured as described in examples 4 and 5. DNA polymerase, for example, Klenow fragment, and deoxynucleotide triphosphates are added in appropriate buffer with random hexamer oligonucleotides to bring about non-specific double-stranded DNA syntheses. The in vitro synthesized DNA product is then made single stranded by heat treatment (e.g., 100°C for three minutes), or its equivalent, and additional DNA polymerase is then added to replace that rendered inactive by the heat treatment. Further in vitro DNA replication then is allowed to occur. The heat treatment and polymerization reactions are repeated about 10 times to produce an approximately 1,000-fold increase in the level of target DNA. The replicated DNA is denatured in vitro using heat or alkali and then captured and detected as described previously.

In this example, rRNA or RNA transcribed from target DNA is purified using a capture probe, described above. The hybrid duplex is then denatured and single stranded nucleic acids are then replicated non-specifically using Q β replicase (methods in Enzymology (1979) 60:628. This replicase replicated both messenger RNA and ribosomal RNA non-specifically under the conditions described by Blumental, Proc. Natl. Acad. Sci. U.S.A. 77:2601, 1908. Because the replication product is a template for the enzyme, the RNA is replicated exponentially.

66

tion is capable of variation and modification and,
therefore, should not be limited to the precise details
set forth, but should include such changes and altera-
tions that fall within the purview of the following
5 claims.

10

15

20

25

30

35

16E050" 0808E280

61

We Claim:

1. A method of amplification of target polynucleotide molecule potentially contained in a sample with non-target polynucleotides comprising the steps of:
- 5 a. contacting the sample potentially containing the target with a first support capable of specifically associating with the target under binding conditions.
- 10 b. separating said first support from the remaining sample to form a removal product which in the presence of target includes target;
- c. subjecting said removal product to amplification which in the presence of target forms an amplification product.
- 15 2. The method of Claim 1 wherein said method further includes the step of determining the presence of said amplification product.
- 20 3. The method of Claim 2 wherein said step of determining the presence of said amplification product includes contacting said amplification product with a probe having a label moiety.
- 25 4. The method of Claim 1 wherein said method further includes the step of contacting said amplification product with a second support capable of specifically associating with said amplification product under binding conditions.
5. The method of Claim 1 wherein said first support includes a retrievable support.
- 30 6. The method of Claim 5 wherein said first support is capable of associating with the target through a probe.
7. The method of claim 6 wherein said probe includes a ligand capable of binding to an antiligand associated with said first support.
- 35 8. The method of Claim 6 wherein said probe is coated with recA protein.

44E050-0308E380

9. The method of claim 1 wherein said amplifying step comprises treating said target with a polymerase.

10. The method of claim 9 wherein said polymerase is RNA polymerase, Q β replicase, transcriptase or DNA polymerase.

11. The method of claim 9 wherein said target is DNA and, prior to said step of treating said target with said polymerase, said target is caused to replicate by subjecting said target to DNA polymerase and non-specific oligonucleotide primer.

12. The method of Claim 1 wherein said target is mRNA.

13. A method of amplifying target polynucleotide molecules potentially contained in a sample with non-target polynucleotides, said method comprising:

a) contacting said sample with a first polynucleotide probe under binding conditions, said first probe capable of specifically associating with said target under binding conditions to form a first probe-target complex;

b) substantially separating said first probe from said non-target polynucleotides in said sample to form a removal product which in the presence of target includes the first probe-target complex;

c) subjecting said removal product to amplification to form an amplification product the generation of which is dependent on the presence of target.

14. The method of claim 13 wherein said amplification product is detectable with a labeled probe.

15. The method of claim 13 further comprising the step subjecting said removal product and said amplification product, if present, to contact with a second probe under binding conditions said second probe capable of specifically associating with said amplification product to form a second probe-amplification product complex, said second probe capable of associating or in associa-

00228060 050394
16E050 1608E280

16. The method of claim 15 wherein said further processing includes the steps of contacting said removal product and said amplification product, if present, with a label probe under binding conditions, said label probe capable of specific association with said amplification product under binding conditions, monitoring said amplification product for the presence of the label probe indicating the presence of the target molecule.

18. The method of claim 17 wherein said polymerase
15 is RNA polymerase, Q β polymerase, reverse transcriptase
or DNA polymerase.

20. The method of claim 13 wherein said target polynucleotide is mRNA.

30 a) a first probe capable of binding to a retrievable support and said target under binding conditions;

35 b) a retrievable support capable of forming a substantially homogeneous dispersion within a sample medium and capable of separation therefrom to form a removal product which in the presence of target in the sample includes said target; and

c) amplification reagents adapted to be applied to said removal product.

22. The method of claim 21 wherein said retrievable support includes at least one bead.

23. The method of claim 22 wherein said bead is capable of interacting with a magnetic field.

24. An instrument for performing assays for target polynucleotides comprising :

a reaction chamber adapted for receiving target polynucleotides and non-target polynucleotides in a sample medium and a support capable of a substantially homogeneous dispersion within the sample medium and capable of forming a complex with the target;

means for separating the support from the sample medium to form a removal product, which in the presence of target in the medium includes target, which removal product is substantially free of non-target polynucleotides;

means to amplify said target as part of a removal product to form an amplification product; and

means to detect said amplification product.

440050-0202E230

add
add
B3

As a below named inventor and petitioner in the foregoing petition, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name

I believe I am an original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled **TARGET AND BACKGROUND CAPTURE METHODS WITH AMPLIFICATION FOR AFFINITY ASSAYS**

the specification of which

(check one)

☒ is attached hereto,

☐ was filed on _____ as

Application Serial No. _____

and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, § 1.56(a).

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application

922,155

October 23, 1986

Pending

Application Serial No.

Filing Date

Status (patented, pending, abandoned)

Application Serial No.

Filing Date

Status (patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of first joint inventor

Mark L. Collins

Inventor's signature

Mark L. Collins

Date

Dec 17, 1987

Residence

435 Malden Street, Holden, Massachusetts 01520

Citizenship

United States

MA

Post Office Address

435 Malden Street, Holden, Massachusetts 01520

Full name of second joint inventor

Donald N. Halbert

Inventor's signature

Donald N. Halbert

Date

Dec. 17, 1987

Residence

31 Janock Road, Milford, Massachusetts 01757

Citizenship

United States

MA

Post Office Address

31 Janock Road, Milford, Massachusetts 01757

Full name of third joint inventor

Walter King

Inventor's signature

Walter King

Date

Dec 14, 1987

Residence

Three Fletcher Street, Maynard, Massachusetts 01754

Citizenship

United States

MA

Post Office Address

Three Fletcher Street, Maynard, Massachusetts 01754

*Dr. Jonathan M. Lawrie's signature to the above declaration is contained following page.

fourth joint inventor Jonathan M. Lawrie

signature

Jonathan M. Lawrie

12/17/87

Residence S. Oak Terrace, Milford, Massachusetts 01757

Citizenship United States

Post Office Address S. Oak Terrace, Milford, Massachusetts 01757

00238000-050344

Serial No.: 644,967
Filed: January 22, 1991

Petition and Power of Attorney

To the Commissioner of Patents and Trademarks:

Your petitioners, Mark L. Collins whose
Post Office address is 435 Malden Street
and who is a resident of Holden, County of Worcester
State of Massachusetts; Donald N. Halbert whose
Post Office address is 31 Janock Road
and who is a resident of Milford, County of Worcester
State of Massachusetts, ~~and~~ Walter King whose
Post Office address is Three Fletcher Street
and who is a resident of Maynard, County of Middlesex
State of Massachusetts and Jonathan M. Lawrie whose
Post Office address is S. Oak Terrace,
and who is a resident of Milford, County of Worcester;
State of Massachusetts;

citizens of the United States, pray that Letters Patent be granted to them for the improvement in

TARGET AND BACKGROUND CAPTURE METHODS WITH AMPLIFICATION FOR AFFINITY ASSAYS

set forth in the annexed specification; and they hereby appoint Ralph C. Medhurst Registration No. 20,082; William H. Magidson Registration No. 19,902; Robert J. Wagner Registration No. 24,186; and Anthony J. Janiuk

Registration No. 29,809

their attorneys; the address of each being, Amoco Corporation, Patents and Licensing Department, Mail Code 1907A, 200 East Randolph Drive, P. O. Box 87703, Chicago, Illinois 60680-0703; each of said attorneys to have full power to prosecute this application, to receive the patent, and to transact all business in the Patent and Trademark Office connected therewith, said Ralph C. Medhurst, said William H. Magidson, and said Robert J. Wagner each to have in addition full power of substitution, association and revocation, including the power to revoke the power of attorney of said Anthony J. Janiuk

The Patent and Trademark Office is requested to direct all mail to said Anthony J. Janiuk

and all telephone calls to said Anthony J. Janiuk

at 312- 856-7972